for immediate and lasting results in HYPER-ACIDITY SYNDROME • Gastritis • Flatulence • Dyspensia • Peptic Ulcer (relief within 5-15 minutes)



Sookty bhasma, Kapoor Kachli, Jatamansi, Ganthoda, Khurasani aimo, Kel pan raakh, Vacha, Dhatura leaves (Detoxicated)

Even in severe cases: 3-6 tablets at a time give relief within 5-15 minutes. In Chronic cases: Gives quick & long lasting relief.

SOOKTYN is a formula of selected Ayurvedic drugs, in the over all management of Hyperacidity and allied conditions.

positive advantages of SOOKTYN treatment

• Detoxicates the digestive tract. • Protects stomach and intestine from Irritants and relieves pain. • Exerts action throughout the length of the gut and allows thorough elimination of unabsorbed residues and thus helps in normal peristalsis. • Having liver corrective properties. Helps digestion, assimilation, morning evacuation and relieves gastric irritation. • Helpful in healing of Duodenal and Gastric ulcers. • Exerts beneficial effect on the central regulatory mechanism. • Safe Tranquillizing effect without any harmful effect on G.I. System.

INDICATION

Hyperacidity syndrome, heart-burn, pain after food, acid eructation, nausea, dyspepsia, ptyalism, vomiting.
Loss of appetite, flatulence, gastritis.
during Pregnancy : for heart-burn, nausea, vomiting.

• pre-operative: to control symptoms in duodenal and gastric ulcers. • Drug induced gastritis: e.g. gastritis due to treatment with aspirin, pyrazolones, indomethacin, cortisone, broad spectrum antibiotics etc.

In the Aged, Hospitalised & Bed ridden patients: to help digestion, assimilation and morning evacuation. Children: In flatulence, dyspepsia, gripe symptoms, vomiting and loss of appetite.

In functional dyspepsia & hyperacidity syndrome : to relieve nocturnal pain; pain free intervals are long and in some cases pain completely disappears.

Dyspepsia & hyperacidity syndrome due to: Alcoholic drink, excess of hot or cold drinks e.g. coffee, tea, aerated water;

heavy smoking and tobacco chewing.

Nausea & discomfort: after heavy or disagreeable meals or drinks.

Before, during or after alcoholic drinks: to avoid untoward effects e.g. dyespepsja nausea, vomiting & hangover. Also improves bowel movement and promotes feeling of freshness next-day.

DOSE

2 tabs. thrice a day between or after principal meals.

In severe symptoms: 3-6 tabs. three or four times a day. Then gradually reduce the dose.

In the Aged, Hospitalised or Bed-ridden: 1-2 tabs. thrice a day after meals.

In children : 1/2 to 1 tabs thrice a day. Tablet can be crushed and mixed with milk, fruit juice or water for easy administration.

SOOKTYN[®]available at Chemists in PACKS of 50 & 100 tablets

ALARSIN 205, GIDC, Pandesara, Surat, Gujarat. **H.O.** : Alarsin House, A/32, Road No. 3, M.I.D.C., Andheri (E), Mumbai 400 093.